



SOFT WHEAT STANDARDS 2020-2021

THIS CHART IS INTENDED TO ACT AS A GUIDE ONLY

Issued on 01 August 2020

Varietal Restrictions

Varietal restrictions apply to the grades on this chart. A variety must not be binned above its maximum allowable varietal classification.

ANW1, ANW2, ASWS – Only the variety Rosella. Where these segregations are not available, these varieties may only be received into FED1 or lower grades.

SFE1, SFE2, SGP1, SGP2 - Only the varieties Anlace, Barham, Gazelle, Impala, Orion, QAL2000. Yenda may be received into SFE1, SFE2, SGP1 or SGP2 segregations (subject to Varietal Classification Zones).

Where these segregations are not available, these varieties may only be received into FED1 or lower grades.

These standards are to be applied on individual truck loads and must not be averaged over a number of loads. Segregations indicated on this chart are only available where announced.

Test Code	Binned Grade	NOODLE			SOFT			
		ANW1	ANW2	ASWS	SFE1	SFE2	SGP1	SGP2
MOGR	Moisture Maximum (%)	12.5	12.5	12.5	12.5	12.5	12.5	12.5
PRGR	Protein Minimum (%)	9.5	No Limit	No Limit	No Limit	No Limit	No Limit	10.5
	Protein Maximum (%)	11.5	No Limit	9.5	9.5	10.5	10.5	No Limit
TWT	Test Weight Minimum (kg/hl)	76	72	76	76	76	70	70
SCRN	Screenings (All Material Below the Screen) Maximum (% by weight)	5.0	10	5.0	8.0	8.0	8.0	8.0
	Material passing through a 2.0mm slotted screen after 40 shakes.							
UNM	Unmillable Material Above the Screen Maximum (% by weight)	0.6	0.6	0.6	0.6	0.6	1.2	1.2
	This consists of whiteheads (with grains removed), chaff, backbone, Wild Radish pods, Milk Thistle pods, Skeleton Weed Flowers (Seed Heads), other seedpods and other light material which remains above the 2.00mm screen after a sample of grain is subjected to the screening process. It excludes contaminants for which tolerances have been stated in these Standards.							
FALL	Falling Number Minimum (seconds)	300	300	300	300	300	250	250
Defective Grains – Maximum tolerance as per method specified. Note: NIL tolerance applies to the entire load.								
Defective definitions are to be read in conjunction with the photo in the Visual Recognition Standards Guide which depicts the minimum standard for a grain to be classified as defective.								
SPRO	Sprouted Maximum (% by count per 300 Grain Sample)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
	Sprouted kernels are those in which the covering of the germ is split and any further advanced stage of growth to the extent of the germ having grown or shot. Note: The NIL tolerance for sprouted grains does not apply if a Falling Number analysis is conducted.							
STAN	Stained Maximum (% by count per 300 Grain Sample)	5% (15 grains)	15% (45 grains)	5% (15 grains)	10% (30 grains)	15% (45 grains)	15% (45 grains)	15% (45 grains)
	A distinct dark brown to black discolouration on the germ end that, in severe cases, may progress to other parts of the grain such as the crease. Grains are commonly referred to as "black point" or "black tip". Includes any staining beyond the minimum and up to 50% of the entire grain surface. For staining level greater than 50%, refer to Field Fungi. Also includes adherence of contaminants such as soil, dust, plant parts and other material.							
PFUN	Pink Stained Maximum (% by count per 300 Grain Sample)	2% (6 grains)	5% (15 grains)	2% (6 grains)	2% (6 grains)	5% (15 grains)	5% (15 grains)	5% (15 grains)
	Grains with distinct pink discolouration. Grains that are pink but also contain a white to light grey fungal like discolouration over more than approximately 50% of the seed coat surface are to be classified as "White Grain Disorder / Head Scab".							
WGDH	White Grain Disorder/Head Scab Maximum (% by count per 300 Grain Sample)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)
	Grains appear white to light grey but may also contain a pink discolouration. Discolouration must be over more than approximately 50% of the seed coat surface. If the discolouration is less than approximately 50% of the seed coat surface, grains may be classified as sound. Grains may also appear flaky with a white discolouration or resemble tombstones.							
GREE	Dry Green or Sappy Maximum (% by count per 300 Grain Sample)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	5% (15 grains)	3% (9 grains)	5% (15 grains)
	Distorted (% by count per 300 Grain Sample)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	5% (15 grains)	3% (9 grains)	5% (15 grains)
DIST	Grains generally have the appearance of full sized kernels with little or no structure on both dorsal sides of the grain, and are typically grey to blue in colour. Does not include pinched grain.							
	Insect Damaged Maximum (% by count per 300 Grain Sample)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	3% (9 grains)	3% (9 grains)
ARTD	Over-Dried Damaged Maximum (% by count per 300 Grain Sample)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
	Field Fungi Maximum (count per half litre)	10	10	10	10	10	10	10
FFUN	Individual kernels where more than half (50%) of the seed coat is discoloured. Discolouration can vary from dark grey, brown to black in colour. Grains that are approximately 50% or less discoloured are to be classified as Stained. Does not include Severely Damaged.							
	Severely Damaged (count per half litre above 2.0mm screen)	1	1	1	1	1	1	1
SEVE	Mould, heat damaged/burnt, or other serious visual defects. Grains that have become severely discoloured. Grains appear reddish brown, dark brown, or in severe cases, blackened. Does not include Field Fungi.							
	All Smuts (except Loose Smut) Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
SMUT	Includes: Ball Smut, Stinking Smut, Bunt, Covered Smut.							
	Foreign Seed Contaminants – Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per half litre.							
Except TYPE(1) in which the maximum applies on an individual seed basis per half litre.								
WS1	TYPE(1): Colocynthis, Double Gees/Spiny Emex/Three Corned Jack, Jute, Long Head Poppy, Mexican Poppy, Opium Poppy, Field Poppy, Horned Poppy, Wild Poppy, New Zealand Spinach, Parthenium Weed** (QLD only)	8*	8*	8*	8*	8*	8*	8*
WS2	TYPE(2): Castor Oil Plant, Coriander, Crow Garlic/ Wild Garlic, Darling Pea, Parthenium Weed** (NSW/VIC/SA), Peanut seeds and pods, Ragweed, Rattlepods, Starburr, St. John's Wort	NIL	NIL	NIL	NIL	NIL	NIL	NIL
WS3A	TYPE(3a): Bathurst Burr, Bellvine, Branched Broomrape, Bulls Head/Caltrop/Cats Head, Cape Tulip, Cottonseed, Dodder, Noogoora Burr, Thornapple	2	2	2	2	2	2	2
WS3B	TYPE(3b): Vetch (Tare), Vetch (Commercial)	4	4	4	4	4	4	4
WS3C	TYPE(3c): Heliotrope (Blue), Heliotrope (Common) ***Note: included in this Type are tolerances for seeds or pods.	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds
WS4	TYPE(4): Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/King Island Mellilot *** (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20	20	20	20	20	20	20
WS5	TYPE(5): Knapweed (Creeping/Russian), Sesbania Pea, Patterson's Curse/ Salvation Jane	40	40	40	40	40	40	40
WS6	TYPE(6): Colombus Grass, Johnson Grass, Saffron Thistle	10	50	10	10	10	10	10
WS7A	TYPE(7a): Adzuki Beans, Broad Beans, Chickpeas, Corn (Maize), Cowpea, Faba Beans, Lentils, Lupins, Peas (Field), Medic Pods, Safflower, Soybean, Sunflower and any other seeds or pods greater than 5mm in diameter. Includes Onion Weed Pods regardless of size	1	10	1	1	1	1	1
WS7B	TYPE(7b): Barley (2 & 6 row), Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Red/ Spring Feed Wheats, Oats (Black/Wild)*, Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Grain), Triticale, Turnip Weed Pods and any other Foreign Seeds not specified in Types 1-7(a), in SFS or in Unmillable Material Above the Screen that remain above the 2.0mm screen following the Screenings process.	50	150	50	50	50	50	50
SFS	Small Foreign Seeds maximum (% by weight)	0.6%	1.2%	0.6%	0.6%	0.6%	0.6%	0.6%
All Foreign Seeds not specified in Types 1-7(b) that fall below the 2.0mm screen during the Screenings process								
* Individual Seed Basis ** Parthenium Weed is a NIL tolerance in NSW/VIC/SA *** Hexham Scent is only acceptable if no tainting odour is present **** Heliotrope pods must be opened and the seeds counted								
Other Contaminants - Maximum tolerance as per method specified. Note: NIL tolerance applies to the entire load.								
PICK	Pickling Compounds Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
CHEM	Chemicals Not Approved for Wheat Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
ERGR	Ryegrass Ergot Maximum (length in cm when pieces are aligned per half litre)	2cm	2cm	2cm	2cm	2cm	2cm	2cm
ERGW	Cereal Ergot Maximum (count per half litre)	1	1	1	1	1	1	1
LIVE	Stored Grain Insects and Pea Weevils: Live Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
INLG	Field Insects – Large, dead or alive Maximum (count per half litre)	3	3	3	3	3	3	3
	Includes: Desiantha Weevil, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, grasshoppers and wood bugs, whole or parts thereof. These are insect contaminants of grain that do not cause damage to stored grains.							
INSM	Field Insects – Small, dead or alive Maximum (count per half litre)	10	10	10	10	10	10	10
	Includes: all species of aphid, minute mould beetle, mites & stored grain insects (dead only). These are insect contaminants of grain that do not cause damage to stored grains.							
SNAL	Snails (Live or Dead) Maximum (count per half litre)	1	1	1	1	1	1	1
EAR	Earcrocodile Maximum (count per half litre)	10	10	10	10	10	10	10
LSMT	Loose Smut Maximum (count per half litre)	3	3	3	3	3	3	3
SAND	Sand Maximum (count per half litre)	20	20	20	20	20	20	20
EART	Earth Maximum (count per half litre) Defined as a clod of dirt, being 5mm or less in diameter.	1	3	1	1	1	1	1
STON	Stones (g per 2.5L above 2.0mm screen) Maximum weight of all stones retained above a 2.0mm screen per 2.5L	4	4	4	4	4	4	4
OBJM	Objectionable Material Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL
	Presence of meat meal, blood meal, fish meal, poultry offal meal or other animal proteins. Sticks/Stubble (>3cm in length and/or 1cm in diameter), glass, concrete, metal, fertiliser, animal excreta, animal carcasses, tainting agents or any other commercially unacceptable contaminant, odour or taste greater than the tolerance or specifications as allowed in these standards.							
OFM	Other Foreign Material Maximum (% by weight)	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%
	Includes fine material (dust and minerals), pieces of snail shell (<half), pieces of stored grain insects, other non-vegetative material not listed.							